

Production, isolation and necrotrophic activity of selected *Parastagonospora nodorum* proteinaceous toxins.

Piotr Ochodzki, Jakub Walczewski, Edward Arseniuk

Department of Plant Pathology, Plant Breeding and Acclimatization Institute - NRI, Radzików, 05-870 Błonie, Poland



One of the major pathogens of triticale and wheat is *Parastagonospora nodorum* (*Stagonospora nodorum*). Infection causes necrotic spots on leaves in which they are growing, and can spread to the same plant, as well as to neighboring plants. The presence of this pathogen during high humidity and temperature can lead to a serious reduction in the yield by reducing the photosynthetically active surface of the leaf and damage to the chaff. In the last few years there have been reports of production by *S. nodorum* protein toxins, which play a key role in the induction of necrotic changes in infected tissues of the host. These toxins affect the specific host genes. The positive interaction of the toxin with the products of dominant form of the gene leads to the induction of necrosis, and in the presence of only a recessive form is the insensitivity to a particular toxin. Experiments conducted under both laboratory and field conditions confirmed that the toxins are a major factor in the development of leaf and chaff septoria on wheat. There were no negative effects of the elimination of the dominant allele of susceptibility to the toxin, while the increase in resistance was found, which positively corresponds with the research Oliver and others (2014). Until now seven different proteinaceous toxins have been described. The aim of the studies was production and purification of necrotrophic effectors in amount which allowed to conduct studies on susceptibility of wheat and triticale genotypes.

MATERIAL AND METHODS

Toxin-producing isolates were grown on liquid medium at pH=5.7, containing sodium acetate, ammonium nitrate, magnesium sulfate, potassium dihydrogen phosphate, sucrose and yeast extract on shaker in room temperature, in darkness during 3 weeks. Culture media were filtrated on blotting paper and later on 0,45 µm filter. Filtrates were dialyzed against 20 mM acetate buffer, pH=5.0 to remove LMW compounds (cut-off MW<3 kDa). Particles larger than 3 kDa were concentrated on centrifugal filter and dissolved with 20 mM acetate buffer, pH=5.0. Partially purified extract was fractionated on HiTrap SP XL 5-mL column by elution with acetate buffers pH 5.0 with increasing ionic strength from 50 to 300 mM of NaCl. The obtained fractions were tested for necrotoxicity on the leaves of selected wheat genotype. Fractions with necrotrophic activity were combined and further separated on 80 cm Superdex 75 chromatography column with 20 mM acetate buffer. Collected fractions were tested for necrotrophic activity and active fractions were pooled. SDS-PAGE method was used for evaluation of purity and mass size of active proteinaceous compounds.

Collected fractions and starting extract were infiltrated by syringe to the second leaves of wheat and triticale genotypes placed on agar with 50ppm benzimidazole. After six days of 16/8 photoperiod in room temperature leaves were checked. Active fractions were combined and leaves of wheat and triticale genotypes were infiltrated.




Figure 1. Results of infiltration of set of segregating wheat lines with fractions of *S. nodorum* isolate 76-40 culture filtrate




Figure 2. Fractionation of crude extract of 76-40 isolate on HiTrap SP XL 5 mL column. * - fractions with necrotic activity




Figure 3. Chromatogram of SEC (Superdex G 75 column) of semipurified extract of isolate No76-40

RESULTS

Isolates Sn76-40 and SN 16-8-03 which produced Tox5 only were selected. Resulting culture filtrate were tested for necrotic activity (Fig.1). Semipurified filtrates fractionated on ion exchange (IEC) (Fig.2) showed 2 fractions with necrotic activity (Fig.2, Fig. 4). Size exclusion chromatography (SEC) of pooled IEC fractions (Fig.3) revealed necrotic activity in fractions #24 and #25. Toxicity of separated fractions tested on the leaves of selected wheat genotype, sensitive to crude culture filtrate confirmed necrotic activity of fractions. The molecular weight of Tox-5 was estimated at approximately 18 kDa (Fig.5.). Susceptibility to Tox5 toxin of ca. 800 wheat and triticale genotypes is evaluated in field experiment.




Figure 4. Results of infiltration of wheat LP 749-29 leaves with fractions of *S. nodorum* isolate 76-40 culture filtrate purified on HiTrap SP XL. * - necrotic active fractions (see Fig.3)




Figure 5. Results of SDS-PAGE of SEC fractions isolate 76-40 (A) and isolate 16-8-03, Fractions #24 and 25 revealed necrotic activity